



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of Nobuhiro ISHIDA et al.

Application Serial No. 10/507,129

FOR: METHOD OF CONTROLLING ETHANOL PRODUCTION AND MASS
PRODUCTION OF LACTIC ACID AND TRANSFORMANT THEREFOR

VERIFICATION OF TRANSLATION

Commissioner for Patents

P.O. Box 1450

Alexandria, VA 22313-1450

Sir:

I, Wakana MITAMURA, a citizen of Japan residing at c/o OKADA PATENT & TRADEMARK OFFICE, P.C., Nagoya Chamber of Commerce & Industry Building, 10-19, Sakae 2-chome, Naka-ku, Nagoya-shi, Aichi 460-0008 JAPAN declare that:

- 1) I can read and understand both Japanese and English;
- 2) I have read the Japanese Patent Application No. 2002-065880;
- 3) the attached English translation is a true and accurate translation of the Patent Application to the best of my knowledge and belief; and
- 4) all statements made of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 USC 1001, and that such false statements may jeopardize the validity of the application or any patent issuing thereon.

Nov. 8, 2007

Date November 8, 2007

W. Mitamura
Wakana MITAMURA

7

[Name of Document]	Patent Application
[Reference Number]	010770
[Date of Filing]	March 11, 2002
[Destination]	Commissioner, Patent Office
[International Classification]	C12N
[Inventor]	
[Address]	c/o KABUSHIKI KAISHA TOYOTA CHUO KENKYUSHO 41-1, Aza Yokomichi, Oaza Nagakute, Nagakute-cho, Aichi-gun, Aichi-ken, JAPAN Nobuhiro ISHIDA
[Name]	
[Inventor]	
[Address]	c/o KABUSHIKI KAISHA TOYOTA CHUO KENKYUSHO 41-1, Aza Yokomichi, Oaza Nagakute, Nagakute-cho, Aichi-gun, Aichi-ken, JAPAN Masana HIRAI
[Name]	
[Inventor]	
[Address]	c/o KABUSHIKI KAISHA TOYOTA CHUO KENKYUSHO 41-1, Aza Yokomichi, Oaza Nagakute, Nagakute-cho, Aichi-gun, Aichi-ken, JAPAN Takao IMAEDA
[Name]	
[Inventor]	
[Address]	c/o KABUSHIKI KAISHA TOYOTA CHUO KENKYUSHO 41-1, Aza Yokomichi, Oaza Nagakute, Nagakute-cho, Aichi-gun, Aichi-ken, JAPAN Chikara MIYAZAKI
[Name]	
[Inventor]	
[Address]	c/o KABUSHIKI KAISHA TOYOTA CHUO KENKYUSHO 41-1, Aza Yokomichi, Oaza Nagakute, Nagakute-cho, Aichi-gun, Aichi-ken, JAPAN Kenro TOKUHIRO
[Name]	
[Inventor]	
[Address]	c/o TOYOTA JIDOSHA KABUSHIKI KAISHA 1, Toyota-cho, Toyota-shi, Aichi-ken, JAPAN Satoshi SAITO
[Name]	
[Inventor]	
[Address]	c/o TOYOTA JIDOSHA KABUSHIKI KAISHA 1, Toyota-cho, Toyota-shi, Aichi-ken, JAPAN Osamu SAOTOME
[Name]	
[Inventor]	
[Address]	c/o TOYOTA JIDOSHA KABUSHIKI KAISHA 1, Toyota-cho, Toyota-shi, Aichi-ken, JAPAN Noriko YASUTANI
[Name]	
[Inventor]	
[Address]	c/o TOYOTA JIDOSHA KABUSHIKI KAISHA 1, Toyota-cho, Toyota-shi, Aichi-ken, JAPAN Yasuo MATSUO
[Name]	
[Applicant]	
[Identification Number]	000003609
[Name]	KABUSHIKI KAISHA TOYOTA CHUO KENKYUSHO

[Applicant]

[Identification Number] 000003207

[Name] TOYOTA JIDOSHA KABUSHIKI KAISHA

[Attorney]

[Identification Number] 100064344

[Patent Attorney]

[Name] Hidehiko OKADA

[Telephone Number] (052) 221-6141

[Designated Attorney]

[Identification Number] 100087907

[Patent Attorney]

[Name] Tetsuo FUKUTA

[Designated Attorney]

[Identification Number] 100105728

[Patent Attorney]

[Name] Atsuko NAKAMURA

[Indication of Fees]

[Number of Advance Payment Ledger] 002875

[Amount of Payment] 21,000

[List of Accompanying Documents]

[Name of Document]	Specification	1
--------------------	---------------	---

[Name of Document]	Set of Drawings	1
--------------------	-----------------	---

[Name of Document]	Abstract	1
--------------------	----------	---

[Name of Document]

Specification

[Title of Invention] TRANSFORMANT HAVING HIGH- LACTIC ACID PRODUCTIVITY
AND USAGE OF THE SAME

[Claims]

[Claim 1] A transformant into which has been incorporated a DNA for coding a foreign protein having lactate dehydrogenase activity and provided with pyruvic acid substrate affinity that equals or exceeds the pyruvic acid substrate affinity of the pyruvate decarboxylase inherent in the host organism.

[Claim 2] The transformant according to Claim 1, wherein the DNA for coding the aforementioned foreign protein has been controllably incorporated by the promoter of the pyruvate decarboxylase 1 gene on the host chromosome or by a homologue of said promoter that replaces said promoter.

[Claim 3] The transformant according to Claim 1 or Claim 2, wherein the aforementioned host organism belongs to the *Saccaromyces* family.

[Claim 4] The transformant according to Claim 1 or Claim 2, wherein the aforementioned host organism is *Saccaromyces cerevisiae*.

[Claim 5] The transformant described in any one of the Claims 1-4, wherein said foreign protein is a bovine-derived lactate dehydrogenase or its homologue.

[Claim 6] A transformant into which has been incorporated DNA for coding a foreign protein having lactate dehydrogenase activity and provided with pyruvic acid substrate affinity that equals or exceeds the pyruvic acid substrate affinity of the pyruvate decarboxylase inherent in the host organism, wherein the DNA for coding the aforementioned foreign protein has been controllably incorporated by the promoter of the pyruvate decarboxylase 1 gene on the host chromosome or by a homologue of said promoter that replaces said promoter.

[Claim 7] The transformant according to Claim 6, wherein the aforementioned promoter on the host chromosome is the promoter of the pyruvate decarboxylase 1 gene.

[Claim 8] The transformant according to Claim 6 or Claim 7, wherein the aforementioned host organism belongs to the *Saccaromyces* family.

[Claim 9] The transformant according to Claim 6 or Claim 7, wherein the aforementioned host organism is *Saccaromyces cerevisiae*.

[Claim 10] The transformant described in any one of the Claims 6-9, wherein said foreign protein is a bovine-derived lactate dehydrogenase or its homologue.

[Claim 11] A transformant into which the DNA for coding the bovine-derived lactate dehydrogenase or its homologue has been controllably incorporated by the promoter of the pyruvate decarboxylase 1 gene on the host chromosome of the *Saccaromyces* family or by a homologue of said promoter that replaces said promoter.

[Claim 12] The transformant according to Claim 10, wherein the aforementioned host is *Saccaromyces cerevisiae*.

[Claim 13] A lactic acid manufacturing method provided with a process for culturing the transformant described in any one of the Claims 1-12, and

a process for separating lactic acid from the cultured product obtained in the aforementioned process.

[Detailed Description of the Invention]

[0001]

[Technical Field of the Invention]

The present invention relates to ethanol production control and a high-productivity technology for lactic acid, and more particularly to a high expression system suitable to lactic acid production using yeast.

[0002]

[Prior Art]

Due to advances in recombinant DNA technology, technologies have been developed that obtain the target gene product by making a foreign gene express itself in a host such as a microbe, mold, animal, plant, or insect, and growing the gene's transformant. For example, by culturing yeast or the like, it is possible to produce a large volume of target gene product through fermentation production.

[0003]

There have been several attempts to produce L-lactic acid using yeast. There have also been attempts to produce L-lactic acid by incorporating a bovine lactate dehydrogenase (LDH) gene into *Saccaromyces cerevisiae* (Eri Adhi et al., "Modification of metabolic pathway of *Saccaromyces cerevisiae* by the expression of lactate dehydrogenase and deletion of pyruvate genes for the lactic acid fermentation at low pH value," *J. Ferment. Bioeng.* Vol.86, No.3, 284-289, 1988; Danio Porro et al., "Development of metabolically engineered *Saccaromyces cerevisiae* cells for the production of lactic acid," *Biotechnol. Prog.* Vol.11, 294-298, 1995; Kohyo (Japanese unexamined patent publication) No. 2001-516584). However, high-volume production of L-lactic acid was not observed in either of these reports.

[0004]

It is known that the pyruvate decarboxylase in *Saccaromyces cerevisiae* has 1 to 1.5 mM substrate affinity (Km value) with respect to the pyruvic acid and multiple isozymes. In ordinary yeast, pyruvate decarboxylase 1 is functioning. However, if this protein does not express itself due to the destruction of the main gene, etc., pyruvate decarboxylase 5 functions, thus maintaining ethanol production ("Autoregulation of yeast pyruvate decarboxylase gene expression requires the enzyme but not its catalytic activity," Ines Eberhardt, Hakan Cederberg, Haijuan Li, Stephen Koning, Frank Jordan and Stephen Hohmann, *Eur. J. Biochem.* Vol. 262, 191-201, 1999).

[0005]

[Problems to be Solved by the Invention]

As explained above, a technology for producing L-lactic acid in high volume inside *Saccaromyces cerevisiae* has not previously been perfected.

One of the objectives of the present invention is to provide a technology to stably produce lactic acid in high volume inside a host organism having a pyruvate decarboxylase gene, such as *Saccaromyces cerevisiae*.

[0006]

The inventors of the present invention conducted research focusing on the lactate dehydrogenase (LDH) that is expressed during the production of lactic acid inside *Saccaromyces cerevisiae*, and perfected a following invention:

More specifically, a following means can be provided according to the present invention.

[0007]

(1) A transformant into which has been incorporated a DNA for coding a foreign protein having lactate dehydrogenase activity and provided with pyruvic acid substrate affinity that equals or exceeds the pyruvic acid substrate affinity of the pyruvate decarboxylase inherent in the host organism.

(2) The transformant as described in (1), wherein the DNA for coding the aforementioned foreign protein has been controllably incorporated by the promoter of the pyruvate decarboxylase 1 gene on the host chromosome or by a homologue of said promoter that replaces said promoter.

(3) The transformant as described in (1) or (2), wherein the aforementioned host organism belongs to the *Saccaromyces* family.

(4) The transformant according to (1) or (2), wherein the aforementioned host organism is *Saccaromyces cerevisiae*.

(5) The transformant described in any one of (1)-(4), wherein said foreign protein is a bovine-derived lactate dehydrogenase or its homologue.

(6) A transformant into which has been incorporated DNA for coding a foreign protein having lactate dehydrogenase activity and provided with pyruvic acid substrate affinity that equals or exceeds the pyruvic acid substrate affinity of the pyruvate decarboxylase inherent in the host organism, wherein

the DNA for coding the aforementioned foreign protein has been controllably incorporated by the promoter of the pyruvate decarboxylase 1 gene on the host chromosome or by a homologue of said promoter that replaces said promoter.

(7) The transformant as described in (6), wherein the aforementioned promoter on the host chromosome is the promoter of the pyruvate decarboxylase 1 gene.

(8) The transformant as described in (6) or (7), wherein the aforementioned host organism belongs to the *Saccaromyces* family.

(9) The transformant according to (6) or (7), wherein the aforementioned host organism is *Saccaromyces cerevisiae*.

(10) The transformant described in any one of (6)-(9), wherein said foreign protein is a bovine-derived lactate dehydrogenase or its homologue.

(11) A transformant into which the DNA for coding the bovine-derived lactate dehydrogenase or its homologue has been controllably incorporated by the promoter of the pyruvate decarboxylase 1 gene on the host chromosome of the *Saccaromyces* family or by a homologue of said promoter that replaces said promoter.

(12) The transformant according to (10), wherein the aforementioned host is *Saccaromyces cerevisiae*.

(13) A lactic acid manufacturing method provided with a process for culturing the transformant described in any one of the (1)-(12), and

a process for separating lactic acid from the cultured product obtained in the aforementioned process.

[0008]

According to the present transformant, as a DNA for coding a foreign protein having lactate dehydrogenase activity and provided with pyruvic acid substrate affinity that equals or exceeds the pyruvic acid substrate affinity of the pyruvate decarboxylase inherent in the host organism, is incorporated into the transformant to express the foreign protein.

Pyruvic acid is the common substrate for pyruvate decarboxylase, which leads to ethanol production; lactate dehydrogenase, which acts as a catalyst for lactic acid production; and the like.

In this transformant, lactate dehydrogenase has a higher level of substrate affinity to pyruvic acid than pyruvate decarboxylase; therefore, the ethanol production catalyzed by pyruvate decarboxylase is suppressed and the lactic acid production by the foreign lactate dehydrogenase is promoted. In this way, it is possible to increase lactic acid production by the transformant.

[0009]

[Preferred embodiments of the invention]

Embodiments of the present invention are explained in detail below.

(Transformant)

The foreign protein that is expressed in the transformant has lactate dehydrogenase (LDH) activity, and, the transformant has pyruvic acid substrate affinity that equals or exceeds the pyruvic acid substrate affinity of this enzyme in its relationship with the pyruvate decarboxylase inherent in the host organism that is to be transformed.

The foreign protein that is expressed in the transformant has lactate dehydrogenase (LDH) activity, and in its relationship with the pyruvate decarboxylase inherent in the host organism that is to be transformed, has pyruvic acid substrate affinity that equals or exceeds the pyruvic acid substrate affinity of this enzyme.

Here, LDH is known as an enzyme that acts as a medium for a reaction that produces lactic acid from pyruvic acid in the glycolytic system of an organism such as yeast. Here, lactic acid includes L(+)-lactic acid and D(-)-lactic acid, but would preferably be L(+)-lactic acid. L(+)-lactic is produced by L(+)-LDH and D(-)-lactic acid is produced by D(-)-LDH.

[0010]

LDH can be used as the foreign protein in the present invention. Various types of homologues exist depending on the type of organism, or even inside [single] organisms. In the present invention, the protein having LDH activity includes natural LDH as well as LDH that is artificially synthesized through chemical synthesis or genetic engineering.

The LDH should preferably be derived from a eucaryotic microbe such as yeast; and more preferably from a higher eucaryote, such as a plant, animal, or insect; and even more preferably from even higher eucaryotes, including mammals such as bovines. Bovine-derived LDH is the most preferable. An example of a bovine-derived LDH is the protein consisting of the amino acid sequence shown in sequence number 1.

Bovine-derived LDH that is derived from a muscle, a heart, or the like, is available and can be used. For the enzyme number, EC 1.1.1.27 can be used.

[0011]

Furthermore, the foreign protein in the present invention includes homologues of these types of LDH. LDH homologues include proteins which has LDH activity, with one or several amino acids in an amino acid sequence of a naturally derived LDH replaced, void, inserted, and/or added; and proteins which also has LDH activity that are at least 70%, and more preferably at least 80%, homologous in their amino acid sequence to a naturally derived LDH.

Note that the number of variations within the amino acid sequence is not limited as long as the original protein function can be maintained, but should preferably be within 70% of the total number of amino acids, more preferably within 30%, and most preferably within 20%.

For example, a desirable homologue would be a protein with one or several amino acids in the amino acid sequence shown in sequence number 1 replaced, void, inserted, and/or added, and which has LDH activity; or a protein that is at least 70%, and more preferably at least 80%, homologous in its amino acid sequence to the amino acid sequence shown in sequence number 1, and which also has LDH activity.

Note that the homology in the sequence can be determined using a homology search tool such as one available from BLAST (<http://blast.genome.ad.jp/>), or FASTA (<http://fasta.genome.ad.jp/SIT/FASTA.html>), etc.

[0012]

Modification of amino acid sequence can be achieved by incorporating a variation that could be a replacement, void, an insertion, and/or an addition as needed into the amino acid sequence that is to be modified, using a site-specific dislocation incorporation method (Current Protocols I Molecular Biology edit. Ausubel et al., (1987) Publish. John Wiley & Sons Selection 8.1-8.5), etc. Such a modification is not limited to artificial incorporation or synthesis of a variation, and includes those generated by artificial variation processes as well as those generated by amino acid variations in the natural world.

[0013]

The foreign protein used in the present invention should preferably have high substrate affinity to pyruvic acid. Here, substrate affinity means the Km value in the Michaelis-Menten equation as shown in Formula (1).


Mathematical expression 1

$$V_0 = k_2 E_0 [S] / ([S] + K_m) \quad \dots \dots (1)$$

Note that V_0 is the steady-state initial speed, $[S]$ is the substrate concentration, $[ES]$ is the concentration of a compound consisting of the enzyme and the substrate, E_0 is the total concentration of the enzyme, k_2 is a speed constant for $ES \rightarrow E + S$.

The Michaelis-Menten formula is based on the relationship among the substrate S , the enzyme E , the substrate-enzyme compound ES , and the product P , expressed by Formula (2) below. k_1 is the equilibrium constant when the product P and the enzyme E are generated from the compound ES , and k_1 is the equilibrium constant when the compound ES is dissociated into the enzyme E and the substrate S .

Mathematical expression 2



[0014]

Here, K_m can be defined from the constants using Formula (3) below.

Mathematical expression 3

$$V_0 = k_2 [ES]$$

$$k_1 [E][S] = (k_2 + k_{-1}) [ES]$$

$$\frac{[E][S]}{[ES]} = \frac{k_2 + k_{-1}}{k_1} = K_m \quad \dots \dots (3)$$

Furthermore, as indicated by Formula (4), K_m can be replaced with various concentrations.

Mathematical expression 4

$$K_m = (E_0 - [ES]) [S] / [ES] \quad \dots \dots (4)$$

[0015]

Substrate affinity can be determined by determining the relationship between the substrate concentration $[S]$ and the initial speed V_0 under a constant enzyme concentration, using a graph that shows the substrate concentration on the horizontal axis and the initial speed on the vertical axis, for example. Substrate affinity can also be determined from a Lineweaver-Burk plot.

[0016]

The pyruvic acid substrate affinity of the foreign protein in the present invention, in its relationship with the pyruvate decarboxylase inherent in the host organism, should preferably equal or exceed the pyruvic acid substrate affinity of this enzyme. Substrate affinity should preferably be compared under the same temperature and pH conditions, etc. The temperature and pH conditions can be determined by taking into consideration the environment in which these enzymes catalyze in the host organism. For example, the substrate affinity of these enzymes should preferably be measured under a temperature of between 30 and 37°C and a pH of between 6.0 and 7.5.

[0017]

The pyruvic acid substrate affinity of the foreign protein should preferably be approximately 1.5 mM or less. Because, if the pyruvic acid substrate affinity exceeds 1.5mM, the pyruvate decarboxylase of the host organism tends to react with pyruvic acid. More preferably, the value should be 1 mM or less. Even more preferably, the value should be 0.5 mM or less, and most preferably should be 0.1 mM or less.

Furthermore, the pyruvic acid substrate affinity of the foreign protein should preferably equal or exceed the pyruvic acid substrate affinity of the pyruvate decarboxylase of the host organism and more preferably exceed the pyruvic acid substrate affinity of the pyruvate decarboxylase of the host organism. If the pyruvic acid substrate affinity of the foreign is lower than the pyruvic acid substrate affinity of the pyruvate decarboxylase of the host organism, the pyruvate decarboxylase of the host organism tends to react with pyruvic acid.

[0018]

A DNA for coding such a foreign protein has been incorporated into the present transformant. There is no restriction on the source of this DNA, which can be a cDNA, genome DNA, synthetic DNA, or the like.

Furthermore, this DNA may possess a naturally derived base sequence for coding LDH, or may be a DNA in which part or the whole of such a base sequence is modified and which codes a protein possessing LDH activity. It may also be a DNA possessing a naturally derived base sequence that codes or synthesizes an LDH homologue.

[0019]

The DNA used in the present invention may possess a base sequence that uses the Codon Usage method, which is often used for the host organism to be transformed. For example, the present DNA may possess a base sequence that has been genetically coded using the Codon Usage method in the *Saccharomyces* family, especially *Saccharomyces cerevisiae*.

[0020]

Note that a DNA can be chemically synthesized, or can be synthesized, using the method developed by Fujimoto et al. (Hideya Fujimoto, Synthetic Gene Creation Method, Plant Cell Engineering Series 7, Plant PCR Experiment Protocol, 1997, Shujunsha Co., Ltd., p95-100), which is a known method for synthesizing long-chain DNA.

[0021]

(DNA structure)

By incorporating the DNA for coding the amino acid sequence of the foreign protein into the host organism and letting the protein coded by this DNA express itself, it is possible to produce lactic acid in the host cells.

For transformation, a DNA structure is used that can express a DNA segment comprised of the present DNA inside the host cells. The form of the DNA structure for transformation is not limited in any way, and plasmid (DNA), bacteriophage (DNA), retrotransposon (DNA), or an artificial chromosome (YAC, PAC, BAC, MAC, etc.) can be selected and adopted according to the incorporation mode (inside or outside the gene) for the foreign gene or the type of host cell. Furthermore, there is no structural restriction, such as linear or circular. Therefore, the DNA structure can be provided with the configuration segment of any of these vectors other than that of the present DNA itself. The preferred procaryotic cell vectors, eucaryotic cell vectors, animal cell vectors, and plant cell vectors are well known in the field

[0022]

Note that the plasmid DNA can, for example, be a YCp-type *escherichia-yeast* shuttle vector, such as pRS413, pRS415, pRS416, YCp50, pAUR112, or pAUR123; a YEp-type *escherichia-yeast* shuttle vector, such as pYES32 or YEp13; a YIp-type *escherichia-yeast* shuttle vector, such as pRS403, pRS404, pRS405, pRS406, pAUR101, or pAUR135; a plasmid derived from *escherichia* (a ColE-type plasmid, such as pBR322, pBR325, pUC18, pUC19, pUC119, pTV118N, pTV119N, pBluescript, pHSG298, pHSG396, or pTrc99A; a pIA-type plasmid, such as pACYC177 or pACYC184; a pSC101-type plasmid, such as pMW118, pMW119, pMW218, pMW219, or the like); or a plasmid derived from *Bacillus subtilis* (pUB110, pTP5, etc.). The phage DNA can be a λ phage (Charon4A, Charon21A, EMBL3, EMBL4, λ gt100, gt11, or zap), ϕ X174, M13mp18, or M13mp19, for example. The retrotransposon can be a Ty factor, for example. YAC can be pYACC2, for example.

[0023]

The present DNA structure is produced by severing a fragment containing, for example, the present DNA, using an appropriate limiting enzyme, and inserting the fragment into the limiting enzyme site or multi-cloning site of the vector DNA being used.

The first form of the present DNA structure is provided with a promoter segment that links a DNA segment comprised of the present DNA, so that said DNA segment can express itself. In other words, the present DNA segment is controllably linked by the promoter to the downstream side of the promoter.

[0024]

It is preferable to use yeast to express the present DNA product, i.e., a protein having LDH activity; and therefore, the use of a promoter that expresses itself inside yeast is preferable. For such a promoter, it is preferable to use a pyruvate decarboxylase gene promoter, gal1 promoter, gal10 promoter, heat shock protein promoter, MF α 1 promoter, PH05 promoter, PGK promoter, GAP promoter, ADH promoter, or AOX1 promoter, for example. In particular, a pyruvate decarboxylase 1 gene promoter derived from the *Saccaromyces* family is preferable, and the use of the pyruvate decarboxylase 1 gene promoter derived from *Saccaromyces cerevisiae* is more preferable. This is because these promoters are expressed at high degrees in the ethanol fermentation route of the *Saccaromyces* family (*cerevisiae*). Note that said promoter sequence can be isolated using the PCR breeding method, which uses the genome DNA of the pyruvate decarboxylase 1 gene of a yeast in the *Saccaromyces* family as the mold. The base sequence of said promoter derived from *Saccaromyces cerevisiae* is shown in sequence number 2. For the promoter segment in the present DNA structure, it is possible to use a DNA comprised of the base sequence described in sequence number 2, as well as a DNA that is comprised of this base sequence with one or several bases void, replaced, or added, and which has promoter activity; or a DNA that is hybridized with a DNA formulated from some or all of the sequences in the base sequence shown in sequence number 2 or its complementary strand under stringent conditions and which has promoter activity (in other words, the homologue of said promoter).

[0025]

The second or another form of the present DNA is provided with the present DNA as well as a DNA segment for homologously recombining the host chromosomes. The DNA segment for homologous recombination is a DNA sequence that is homologous to the DNA sequence in the vicinity of the target site in the host chromosome into which the DNA is to be incorporated. At least one, preferably two, DNA segments for homologous recombination should be provided. For example, two DNA segments for homologous recombination should preferably have DNA sequences homologous to the DNA on the upstream and downstream sides of the target site on the chromosome, and the present DNA should preferably be linked between these DNA segments.

[0026]

When incorporating the present DNA into the host chromosome by means of homologous recombination, it is possible to incorporate the present DNA in a state that is controllable by a promoter on the host chromosome. In this case, by incorporating the present DNA, it is possible to simultaneously destroy the endogenous gene that should have been controlled by said promoter and to allow the present DNA, which is foreign, to express itself instead of the endogenous gene. This process is especially useful when said promoter is a highly expressive promoter in the host cell.

[0027]

To create such an expression system on the host chromosome, it is preferable to target a high-expression gene in the host chromosome and to incorporate the present DNA into the downstream side of the promoter that controls this gene, so that the present DNA is controlled by the promoter. If an ethanol-fermenting microbe, such as yeast, is used as the host, it is possible to target a pyruvate decarboxylase gene (particularly, the pyruvate decarboxylase 1 gene) and incorporate a DNA for coding an LDH-active protein under the control of an endogenous pyruvate decarboxylase gene promoter. In this case, it is possible to make the DNA segment for homologous recombination homologous to the sequence in the structural gene region of the LDH of the pyruvate decarboxylase 1 gene or its vicinity (including the sequence near the start codon, the sequence on the upstream region of the start codon, and the sequence inside the structural gene). Preferably, an enzyme in the *Saccaromyces* family (especially *cerevisiae*) should be used as the host, and a DNA structure that targets the pyruvate decarboxylase 1 gene in this host should be used. With such a DNA structure, the destruction of the pyruvate decarboxylase 1 gene and replacement of this structural gene with LDH can be achieved using only a single vector. Because the pyruvate decarboxylase 1 gene is an enzyme that mediates the irreversible reaction from pyruvic acid to acetaldehyde, the destruction of its gene can be expected to suppress the production of ethanol via acetaldehyde, and at the same time promote the production of lactic acid by LDH, which uses pyruvic acid as the substrate.

[0028]

Even the first DNA structure can be made into a DNA structure for homologous recombination by providing it with a DNA segment for homologous recombination with the host chromosome. In the first DNA structure, the promoter segment inside the DNA structure can also be used as the DNA segment for homologous recombination. For example, for the host *Saccaromyces cerevisiae*, a DNA structure that has the promoter in the *Saccaromyces cerevisiae* host chromosome, e.g., the pyruvate decarboxylase 1 gene promoter as a promoter segment comprises a targeting vector that has said gene 1 as the target site. In this case, the first DNA structure should preferably be provided with a sequence homologous to the structural gene region of the pyruvate decarboxylase 1 gene.

[0029]

Note that it is possible to link to the DNA structure, a terminator, and as needed, a cis-element such as an enhancer, a splicing signal, a polyA additional signal, a selective marker, or a ribosome bond sequence (SD sequence). There are no special restrictions on the selective marker, and various types of known marker genes can be used, such as drug-resistant genes or auxotrophic genes. For example, the dihydrofolate reductase gene, ampicillin-resistant gene, neomycin-resistant gene, or the like can be used.

[0030]

(Transformation by DNA structure)

Once a DNA structure has been built, it can be incorporated into an appropriate host cell by means of an appropriate method, such as the transformation method, the transfection method, the bonding method, protoplast fusion, the electroporation method, the lipofection method, the lithium acetate method, the particle gun method, the calcium phosphate precipitation method, the agrobacterium method, the PEG method, or the direct microinjection method. After the incorporation of the DNA structure, the recipient cell is cultured in a selective media.

[0031]

For the host cell, it is possible to use a bacterium such as *Escherichia coli* or *Bacillus subtilis*; a yeast such as *Saccaromyces cerevisiae*, *Saccaromyces pombe*, or *Pichia pastoris*; an insect cell such as sf9 or sf21; an animal cell such as a COS cell or a Chinese hamster ovarian cell (CHO cell); or a plant cell from sweet potato, tobacco, or the like. The host cell should preferably be a microbe, such as yeast, that causes alcoholic fermentation, or an acid-resistant microbe, examples of which include yeasts represented by the *Saccaromyces* family, such as *Saccaromyces cerevisiae*. Specific examples include the *Saccaromyces cerevisiae* IFO2260 strain and YPH strain.

[0032]

In the transformant created by the DNA structure, the structural components of the DNA structure are present on the chromosomes or extrachromosomal elements (including artificial chromosomes). If the DNA structure is maintained outside the chromosome or is incorporated into the chromosome based on random integration, other types of yeast that use pyruvic acid, which is the substrate for LDH, as the substrate, e.g., the pyruvate decarboxylase gene (the pyruvate decarboxylase 1 gene in the case of yeasts belonging to the *Saccaromyces* family), should preferably be knocked out by the targeting vector.

When the DNA structure that is described above and that can achieve homologous recombination is incorporated, a DNA segment, which is the present DNA controllably linked by the desired promoter on the host chromosome, is present on the downstream side of said promoter or a homologue of said promoter that replaces said promoter. In the transformant of a yeast in the *Saccaromyces* family, it is preferable to provide the present DNA on the downstream side of the pyruvate decarboxylase 1 gene promoter on the host chromosome or a homologue of said promoter that replaces said promoter, so that the present DNA can be controlled by said promoter.

Furthermore, a selective marker gene and part of the destroyed structural gene (at the site corresponding to the homologous sequence on the DNA structure) are normally present on the downstream side of the present DNA in a homologous recombinant.

[0033]

Whether or not the present DNA has been incorporated under the desired promoter can be checked using the PCR method or the Southern hybridization method. For example, it is possible to check this by formulating DNA from the transformant, performing PCR with an incorporation site-specific primer, and detecting the expected band from the PCR product in electrophoresis. Alternatively, it is also possible to

perform PCR with a primer that has been marked with a fluorescent dye, etc. These methods are known to those skilled in the art.

[0034]

(Manufacturing of lactic acid)

Culturing the transformant obtained by incorporating the DNA structure produces lactic acid, which is an expression product of the foreign gene, in the cultured product. Lactic acid can then be obtained by separating it from the cultured product. In the present invention, "cultured product" includes the culture supernatant, as well as the cultured cell or microbe, and the crushed form of the cell or microbe.

For culturing the transformant according to the present invention, the culturing conditions can be selected according to the type of transformant. These culturing conditions are known to those skilled in the art.

For the medium for culturing the transformant obtained using a microbe, such as *Escherichia coli* or yeast, as the host, any natural or synthetic medium can be used provided that it contains a carbon source, a nitrogen source, and inorganic salts, etc. that the microbe can utilize, and that it can efficiently culture the transformant. For the carbon source, carbohydrates such as glucose, fructose, sucrose, starch, or cellulose; organic acids such as acetic acid or propionic acid; or alcohols such as ethanol or propanol can be used. For the nitrogen source, ammonium salts of inorganic or organic acids, such as ammonium chloride, ammonium sulfide, ammonium acetate, or ammonium phosphate; other nitrogen-containing compounds; peptone; meat extract; or corn steep liquor can be used. For the inorganics, potassium phosphate, magnesium phosphate, magnesium sulfate, sodium chloride, ferrous sulfate, manganese sulfate, copper sulfate, or calcium carbonate, etc. can be used.

[0035]

Culturing is usually performed under an aerated condition, such as shake culture or aerobic agitation culture, at 30°C for 6 to 24 hours. During culturing, the pH level should preferably be maintained between 2.0 and 6.0. The pH level can be adjusted using an inorganic or organic salt, an alkaline solution, or the like. During culturing, an antibiotic, such as ampicillin or tetracycline, may be added to the medium as needed.

[0036]

For the medium for culturing the transformant obtained using an animal cell as the host, an RPMI1640 medium or a DMEM medium, which are commonly used, or a medium consisting of either of these with bovine fetal serum or the like added to it may be used. Culturing is normally performed under the presence of 5% CO₂ at 37°C for 1 to 30 days. During culturing, an antibiotic such as kanamycin or penicillin may be added to the medium as needed.

[0037]

After culturing is finished, ordinary refining methods can be used in various combination in order to separate a gene product, from the cultured product. For example, if a gene product is produced inside transformed cells, the gene product can be separated from the cells using a regular method, such as ultrasound destruction, grinding, or pressure crushing, to destroy the cells of the bacteria. In this case, protease should be added as needed. If lactic acid is produced in the culture supernatant, solids are removed from this solution through filtering, centrifugal separation, or the like and nucleic acid may be removed through protamine treatment, if necessary.

[0038]

Crudely protein solutions can be obtained by collecting the deposits caused by adding and fractionating ammonium sulfate, alcohol, acetone or the like with respect to these crudely extracted fractions. These protein solutions are treated by means of various chromatography or electrophoresis methods so that a refined enzyme preparation is obtained. For example, a refined objective gene product can be obtained by a gel filtration using Sephadex, Ultra Gel or Bio Gel, an electrocataphoresis method using ion exchange chromatography or polyacrylamide gel, or a fraction method using affinity chromatography or reversed phase chromatography or combining these methods. The amino acid sequence of the refined gene products can be determined by an ordinary amino acid analysis method.

The above described culture methods or refining methods are merely some examples among many various methods and are not limited to these methods.

[0039]

(Embodiments)

Although embodiments of the present invention are described below, they are not intended to specifically limit the scope of the present invention.

[0040]

(Embodiment 1: Design of the DNA sequence of L-lactate dehydrogenase gene)

In order to efficiently produce L-lactate dehydrogenase, which is derived from bovine, a high eucaryote, in the *Saccaromyces cerevisiae* family of yeast, we designed a new, non-naturally occurring gene sequence for the DNA that codes the amino acid sequence of the bovine-derived L-lactate dehydrogenase.

- 1) The codon that is often used in *Saccaromyces cerevisiae* was used.
- 2) Kozaks sequence (ANNATGG) was added straddling the start codon.
- 3) Unstable sequences and repeated sequences in mRNA were eliminated as much as possible.
- 4) The GC content deviation was made uniform over the entire region.
- 5) Measures were taken to ensure that restriction enzyme sites, not suitable to gene cloning, would not be created inside the designed sequence.
- 6) Restriction enzyme *EcoRI*, *XhoI*, and *AflIII* sites, which are useful for the two ends to be incorporated into the chromosome incorporation type vector, were added.

For the codon usage frequency in yeast, the *Saccaromyces cerevisiae* codon usage obtained from the codon usage database (<http://www.kazusa.or.jp/codon/>) is shown in Figure 2. In this figure, the codons that are used frequently for specific amino acids are identified (the underlined codons).

[0041]

A new DNA sequence (999bp) for coding a protein having the LDH activity obtained based on the design guidelines in 2) through 5) above and by applying the frequently used codon in the codon usage shown in Figure 1 (hereafter referred to as "LDHKCB gene") is shown in sequence number 3. Additionally, a DNA sequence (1052bp) that includes the DNA sequence shown in sequence number 3, as well as the upstream side of its start codon and the downstream side of its stop codon (hereafter referred to as "LDHKCB sequence"), is shown in sequence number 4.

[0042]

In the DNA sequence shown in sequence number 3, codons that were different from those used in the original DNA sequence were used in all amino acids except methionine. Note that the newly adopted codons were all frequently used codons from among those shown in Figure 1. Furthermore, Figure 2 shows the result of a computer-based homology analysis of the original bovine-derived LDH gene and the LDHKCB gene. As is clear from Figure 2, it was discovered that a large number of replacements were needed over almost the entire DNA sequence.

[0043]

(Embodiment 2: Complete synthesis of the LDHKCB sequence)

In this embodiment, the method of Fujimoto et al. was used, which is a known method for synthesizing long-chain DNA (Hideya Fujimoto, Synthetic Gene Creation Method, Plant Cell Engineering Series 7, Plant PCR Experiment Protocol, 1997, Shujunsha Co., Ltd., p95-100). The principle behind this method is as follows: oligonucleotide primers of around 100 mer are created so as to have an overlap of between 10 and 12 mer at the 3' end. Then, the missing area is extended using the overlapped region between the oligonucleotide primers, and is amplified through a PCR using the primers at both ends. This operation is sequentially repeated to synthesize the target long-chain DNA. For the PCR amplifier, Gene Amp PCR system 9700 (P.E. Applied Biosystems Inc.) was used.

Specifically, the two kinds of oligonucleotide primers to be linked were mixed first, and a DNA extension reaction was carried out at 96°C for 2 minutes, 68°C for 2 minutes, 54°C for 2 minutes, and 72°C for 30 minutes under the presence of KOD-plus-DNA polymerase (Toyobo). Next, using 1/10 of the sample as a mold, a PCR (polymerase chain reaction) was carried out. In this reaction, the sample was first kept at 96°C for 2 minutes; it was then put through 25 cycles (with each cycle consisting of 96°C for 30 seconds, 55°C for 30 seconds, and 72°C for 90 seconds) under the presence of primers at both ends; and finally was kept at 4°C. For the reaction, the buffer and dNTPmix, etc. that came with the DNA polymerase were used.

A series of overlapping PCRs were carried out according to Figure 3 to create the gene fragments that were ultimately targeted. The DNA sequences of all 28 primers shown in Figure 3 (BA, B01, BB, B02, BC, B03, BD, B04, BE, B05, BF, B06, BG, B07, BH, B08, BI, B09, BJ, B10, BK, B11, BL, B12, BM, B13, BN, and B14) are shown in sequence numbers 5 through 32, respectively. After the base sequence of the synthesized LDHKCB gene was verified, a restriction enzyme process using EcoRI was applied. The sequence was then linked to the pCR2.1 TOPO Vector (Invitrogen), to which an enzyme process using EcoRI had been applied in a similar manner, using a normal method. This vector is referred to as the pBTOPO-LDHKCB vector.

[0044]

(Embodiment 3: Construction of a vector for yeast chromosome incorporation)

A yeast chromosome incorporation-type vector was constructed using the LDHKCB sequence completely synthesized in Embodiment 2. This vector is referred to as pBTRP-PDC1-LDHKCB and its plasmid map is shown in Figure 4.

1. Isolating PDC1P fragments for constructing pBTrp-PDC1-LDHKCB

PDC1P fragments were isolated by means of the PCR amplification method that uses the genome DNA of the *Saccharomyces cerevisiae* YPH strain (Stratagene Corp.) as the mold.

[0045]

The genome DNA of the *Saccharomyces cerevisiae* YPH strain was prepared using the Fast DNA Kit (Bio 101 Inc.), which is a genome preparation kit, and according to the detailed protocol described in the appendix. DNA concentration was measured using the spectrophotometer Ultro spec 3000 (Amersham Pharmacia Biotech Inc.).

[0046]

For the PCR, Pyrobest DNA Polymerase (Takara Shuzo Co., Ltd.), which is said to produce highly precise amplified fragments, was used as the amplification enzyme. A total of 50 µl of reaction solution, containing a 50 ng/sample of the genome DNA of the *Saccharomyces cerevisiae* YPH strain prepared using the aforementioned method, a 50 pmol/sample of the primer DNA, and a 0.2 units/sample of Pyrobest DNA Polymerase, were prepared. The reaction solution was put through DNA amplification using a PCR amplifier system (Gene Amp PCR system 9700 made by P.E. Applied Biosystems Inc.). The reaction conditions for the PCR amplifier were as follows: the solution was first kept at 96°C for 2 minutes; it was then put through 25 cycles (with each cycle consisting of 96°C for 30 seconds, 55°C for 30 seconds, and 72°C for 90 seconds); and finally was kept at 4°C. The PCR-amplified fragments were verified to be amplified gene fragments by means of electrophoresis in 1% TBE agarose gel. Note that a synthetic DNA (Sawaday Technology Co.) was used as the primer DNA for the reaction; the DNA sequence of this primer is described below.

[0047]

- A restriction enzyme BamHI site was added to the PDC1P-LDH-U (31 mer, Tm value of 58.3°C) end.

: ATA TAT GGA TCC GCG TTT ATT TAC CTA TCT C (sequence number 33)

- A restriction enzyme EcoRI site was added to the PDC1P-LDH-D (31 mer, Tm value of 54.4°C) end.

: ATA TAT GAA TTC TTT GAT TGA TTT GAC TGT G (sequence number 34)

[0048]

2. Constructing a recombinant vector containing a promoter and the target gene

Under the control of the promoter sequence for pyruvate decarboxylase 1 gene (PDC1) derived from *Saccharomyces cerevisiae*, a bovine-derived L-lactate dehydrogenase gene (LDH gene) was used as the target gene.

[0049]

The new chromosome incorporation-type vector constructed for constructing the present recombinant vector was named pBTrp-PDC1-LDHKCB. The details of an example of constructing the present vector are described below. An overview of the present embodiment is illustrated in Figures 5 through 8. Note that the procedure for constructing the vector is not limited to that described here.

When constructing the vector, the PDC1 gene promoter fragment (PDC1P) 971bp and the PDC1 gene downstream region fragment (PDC1D) 518bp were isolated using the PCR amplification method that uses the genome DNA of the *Saccharomyces cerevisiae* YPH strain as the mold, as described above. Although the procedure described above was used for PCR amplification, the following primers were used for amplifying the PDC1 gene downstream region fragment:

- A restriction enzyme XhoI site was added to the PDC1D-LDH-U (34 mer, Tm value of 55.3°C) end.
: ATA TAT CTC GAG GCC AGC TAA CTT CTT GGT CGA C (sequence number 35)
- A restriction enzyme ApaI site was added to the PDC1D-LDH-D (31 mer, Tm value of 54. 4°C) end.
: ATA TAT GAA TTC TTT GAT TGA TTT GAC TGT G (sequence number 36)

[0050]

After the PDC1P and PDC1D amplified gene fragments obtained through the aforementioned reactions were each refined through an ethanol precipitation process, the PDC1P amplified fragment and the PDC1D amplified fragment were reacted with restriction enzymes BamHI/EcoRI and XhoI/ApaI, respectively. Note that all of the enzymes used below were made by Takara Shuzo Co., Ltd. Additionally, the series of detailed operations in the ethanol precipitation process and the restriction enzyme processes were based on "Molecular Cloning - A Laboratory Manual, Second Edition" (Maniatis et al., Cold Spring Harbor Laboratory Press, 1989).

[0051]

The series of reaction operations for constructing the vector were carried out according to a generally known DNA subcloning method. That is, the PDC1P fragment, which had been amplified by the aforementioned PCR method and to which a restriction enzyme process had been applied, was linked through a T4 DNA Ligase reaction to the pBluescriptII SK + vector (Toyobo), to which the restriction enzyme BamHI/EcoRI (Takara Shuzo) and the dephosphorylase Alkaline Phosphatase (BAP, Takara Shuzo) had been applied (Figure 5 top row). For the T4 DNA Ligase reaction, the LigaFast Rapid DNA Ligation System (Promega Corporation) was used, following the included detailed protocols.

[0052]

Next, transformation to competent cells was carried out using the solution in which the ligation reaction had taken place. For the competent cells, *Escherichia coli* JM109 strain (Toyobo) was used, following the included detailed protocols. The obtained cultured solution was sprayed onto an LB plate containing the antibiotic ampicillin at the rate of 100 µg/ml, and incubated overnight. The colony that grew was verified with the colony PCR method using the primer DNA of the insert fragment, and with a restriction enzyme process for a plasmid DNA solution prepared through miniprep. The target vector (pBPDC1P vector) was then isolated (Figure 5 middle row).

[0053]

Next, as shown in the middle row in Figure 5, the LDHCKB gene fragment, obtained by processing the pBTOPO-LDHCKB vector constructed by Kabushiki Kaisha Toyota Chuo Kenkyusho. with the restriction enzyme EcoRI and the end-modification enzyme T4 DNA polymerase, was subcloned, using the same operation as described above, into the pPDC1P vector likewise processed with the restriction enzyme EcoRI and the terminal modification enzyme T4 DNA polymerase, to create the pBPDC1P-LDHCKB vector (Figure 5 bottom row).

[0054]

On the other hand, as shown in Figure 6, the LDH gene (derived from *Bifidobacterium longum*) fragment obtained by processing the pYLD1 vector, constructed by Toyota Jidosha Kabushiki Kaisha, with the restriction enzyme EcoRI/AatII and the end-modification enzyme T4 DNA polymerase, was subcloned, using the same operation as described above, into the pBPDC1P vector likewise processed with the restriction enzyme EcoRI and the terminal modification enzyme T4 DNA polymerase, to create the pBPDC1P-LDH1 vector (Figure 6). Note that the aforementioned pYLD1 vector has been incorporated into *Escherichia coli* (name: *E. coli* pYLD1) and has been internationally deposited with the International Patent Organization Depository of the National Institute of Advanced Industrial Science and Technology (Chuo No. 6, 1-1-1 Higashi, Tsukuba-shi, Ibaraki-ken, 305-8566 Japan) under deposit No. FERM BP-7423 in accordance with the Budapest Convention (the original deposit date: October 26, 1999).

[0055]

Next, as shown in Figure 7, this vector was processed with XhoI/ApaI and was linked to the amplified PDC1D fragment, to which a restriction enzyme had been applied in the same manner, to create the pBPDC1P-LDH vector (Figure 7 top row). Lastly, the pBPDC1P-LDHII vector, to which a EcoRV process had been applied, was linked to a Trp marker fragment obtained by applying the AatII/SspI process and the T4 DNA polymerase process to the pRS404 vector (Stratagene Inc.), to construct the pBTrp-PDC1-LDH vector.

[0056]

Next, as shown in Figure 8, the pBPDC1P-LDHKCB vector was processed with restriction enzymes ApaI/EcoRI; separately, the pBTrp-PDC1-LDH vector was processed into fragments containing the Trp marker processes with restriction enzymes ApaI and StuI. The amplified fragments were linked to construct the chromosome incorporation-type pBTrp-PDC1-LDHKCB vector, which is the final construct.

[0057]

In order to verify the constructed chromosome incorporation-type pBTrp-PDC1-LDHKCB vector, base sequence determination was performed. An ABI PRISM 310 Genetic Analyzer (P.E. Applied Biosystems Inc.) was used for analyzing the base sequence, and details such as the method of preparing samples and using the instrument were based on the manual provided with the analyzer. The vector DNA to be used as the sample was prepared with an alkali extraction method. Before using this sample, it was first column-refined using the GFX DNA Purification kit (Amersham Pharmacia Biotech Inc.), and then its DNA concentration was measured with the Ultraspec 3000 spectrophotometer (Amersham Pharmacia Biotech Inc.).

[0058]

(Embodiment 4: Transformation of yeast)

The gene incorporates into yeast according to the method developed by Ito et al. (Ito, H., Y. Fukuda, K. Murata and A. Kimura, Transformation of intact yeast cells treated with alkalinizations, J. Bacteriol. Vol. 153, p163-168). That is, the yeast IFO2260 strain (a strain registered with the Institute of Fermentation, Osaka), from which the tryptophan-synthesizing function had been removed, was cultured in 10 ml YPD medium at 30°C until it reached the logarithmic growth phase. It was then collected and rinsed with TE buffer. Next, 0.5 ml of TE buffer and 0.5 ml of 0.2M lithium acetate were added, and the mixture was shake-cultured at 30°C for 1 hour. Then, the chromosome incorporation-type pBTrp-PDC1-LDHKCB vector constructed using the method in Embodiment 3 was processed with restriction enzymes ApaI and SacI (both made by Takara Shuzo), and was added to the culture.

[0059]

After the present yeast suspension was shake-cultured at 30°C for 30 minutes, 150 µl of 70% polyethylene glycol 4000 (Wako Pure Chemical Industries) was added and the mixture was agitated well. Then, after the mixture was shake-cultured at 30°C for 1 additional hour, a heat shock was applied at 42°C for 5 minutes. The bacteria was then rinsed and suspended in 200 µl of water, and this solution was applied to a tryptophan selective culture medium.

[0060]

The obtained colony was applied to a new tryptophan streak culture medium, and selected strains that had demonstrated stability were checked for gene incorporation using PCR analysis. The genome DNA of the yeast to be used for PCR was prepared by shake-culturing a single colony in 2 ml of YPHD media overnight, collecting the bacteria, adding 50 mM Tris-HCl 500 µl and glass beads (425 - 600 µm, acid washed, SIGMA), and putting the mixture through a vortex at 4°C for 15 minutes. The supernatant of this solution was put through ethanol precipitation and was dissolved in 50 µl of sterilized water. Using 5 µl of the prepared genome DNA as the mold, PCR was performed in 50 µl of reaction solution. EX Taq DNA Polymerase (Takara Shuzo) was used for the DNA amplification enzyme, and the PCR amplifier Gene Amp PCR system 9700 (P.E. Applied Biosystems Inc.) was used. The reaction condition for the PCR amplifier was as follows: the solution was first kept at 96°C for 2 minutes; it was then put through 30 cycles (with each cycle consisting of 96°C for 30 seconds, 55°C for 30 seconds, and 72°C for 90 seconds); and finally was kept at 4°C. The sequences of the primers used were as follows:

LDH-KCB-U : TGG TTG ATG TTA TGG AAG AT (20 mer) (Sequence number 37)

LDH-KCB-D : GAC AAG GTA CAT AAA ACC CAG (21 mer) (Sequence number 38)

[0061]

Those strains that possess a stable tryptophan-synthesizing function and for which PCR was verified under these primers were judged to be transformants into which the LDHKCB gene had been appropriately incorporated. In the present embodiment, three kinds of bacterial strains, namely KCB-27, KCB-210, and KCB-211, were obtained as such transformants. Figure 9 shows the genome chromosome structures of these yeast *Saccharomyces cerevisiae* transformants.

[0062]

(Embodiment 5: Measurement of L-lactic acid production volume in the transformants)

Fermentation experiments were conducted on the three kinds of transformants created. As pre-culturing, the transformants were cultured overnight in a YPD solution medium with 2% glucose concentration. After the bacteria were collected and rinsed, they were planted in a YPD solution medium with 15% glucose concentration such that the bacterial concentration was 1% (0.5 g/50 ml), and were left to ferment at 30°C for several days. This fermented solution was sampled every 24 hours and the volume of L-lactic acid contained was estimated[sic]. For measuring the volume produced, Biosensor BF-4 (Oji Scientific Instruments) was used following the detailed measurement methods described in the user's manual for the instrument.

[0063]

Figure 10 and Table 1 show the measurement results of the L-lactic acid volume and the ethanol volume in the cultured solution on the third day of fermentation with a glucose concentration of 15%. As for the KCB-27 strain, Figure 11 and Table 2 show the trends in L-lactic acid volume and ethanol volume between the 0th and 5th days.

[Table 1]

Strain	Production volume (%)	
	Lactic acid	Ethanol
TDO1	0	7.34
KCB-27	4.92	5.01
KCB-210	4.78	5.14
KCB-211	4.57	5.26

[Table 2]

Strain	Production volume (%)	Number of days passed			
		0	1	3	5
TDO1	Glucose	15	1.84	0	0
	Lactic acid	0	0	0	0
	Ethanol	0	6.69	7.21	7.38
KCB-27	Glucose	15	2.2	0	0
	Lactic acid	0	3.01	4.73	4.92
	Ethanol	0	4.69	5.11	5.06

Even though only a single copy of the LDHKCB gene had been incorporated into each of the transformants into which the LDHKCB gene was incorporated (KCB-27, KCB-210, and KCB-211 strains), production of between 4.5 and 5.0% (45.0 to 50.0 g) of L-lactic acid was confirmed in each 1 L of the culture solution. Meanwhile, ethanol production was 5%, which was approximately 2.5% less than would be expected with the parent strain.

This result clearly indicates an extremely significant increase in the production of L-lactic acid in *Saccharomyces cerevisiae* as compared to the cases reported in the past. Furthermore, this increase in production volume is presumed to be due to the introduction of LDH with substrate affinity equal to or higher than the pyruvic acid substrate affinity of the pyruvate decarboxylase derived from yeast. It is also presumed that introducing LDH under the control of the Pyruvate decarboxylase 1 gene promoter on the chromosome contributed as well.

The above result shows that even a single copy can produce a significant increase in the production volume. Therefore, it can be presumed that the introduction of two or more copies would increase the production volume even further.

[0064]

[Effects of Invention]

The present invention can provide a technology for stably mass-producing lactic acid inside a host organism having the pyruvate decarboxylase gene.

[SEQUENCE LISTING]

SEQUENCE LISTING

<110> Kabushiki Kaisha Toyota Chuo Kenkyusho

Toyota Jidosha Kabushiki Kaisha

<120> Lactate High-expression system and Use thereof

<130> 010770(P01-3840)

<140>

<141>

<160> 39

<170> PatentIn Ver. 2.1

<210> 1

<211> 332

<212> PRT

<213> Bovine

<400> 1

Met Ala Thr Leu Lys Asp Gln Leu Ile Gln Asn Leu Leu Lys Glu Glu

1

5

10

15

His Val Pro Gln Asn Lys Ile Thr Ile Val Gly Val Gly Ala Val Gly

20

25

30

Met Ala Cys Ala Ile Ser Ile Leu Met Lys Asp Leu Ala Asp Glu Val

35

40

45

Ala Leu Val Asp Val Met Glu Asp Lys Leu Lys Gly Glu Met Met Asp

50

55

60

Leu Gln His Gly Ser Leu Phe Leu Arg Thr Pro Lys Ile Val Ser Gly

65

70

75

80

Lys Asp Tyr Asn Val Thr Ala Asn Ser Arg Leu Val Ile Ile Thr Ala

85

90

95

Gly Ala Arg Gln Gln Glu Gly Glu Ser Arg Leu Asn Leu Val Gln Arg

100

105

110

Asn Val Asn Ile Phe Lys Phe Ile Ile Pro Asn Ile Val Lys Tyr Ser

115

120

125

Pro Asn Cys Lys Leu Leu Val Val Ser Asn Pro Val Asp Ile Leu Thr

130

135

140

Tyr Val Ala Trp Lys Ile Ser Gly Phe Pro Lys Asn Arg Val Ile Gly

145

150

155

160

Ser Gly Cys Asn Leu Asp Ser Ala Arg Phe Arg Tyr Leu Met Gly Glu

165

170

175

Arg Leu Gly Val His Pro Leu Ser Cys His Gly Trp Ile Leu Gly Glu

180

185

190

His Gly Asp Ser Ser Val Pro Val Trp Ser Gly Val Asn Val Ala Gly

195

200

205

Val Ser Leu Lys Asn Leu His Pro Glu Leu Gly Thr Asp Ala Asp Lys

210

215

220

Glu Gln Trp Lys Ala Val His Lys Gln Val Val Asp Ser Ala Tyr Glu

225

230

235

240

Val Ile Lys Leu Lys Gly Tyr Thr Ser Trp Ala Ile Gly Leu Ser Val

245

250

255

Ala Asp Leu Ala Glu Ser Ile Met Lys Asn Leu Arg Arg Val His Pro

260

265

270

Ile Ser Thr Met Ile Lys Gly Leu Tyr Gly Ile Lys Glu Asp Val Phe

275

280

285

Leu Ser Val Pro Cys Ile Leu Gly Gln Asn Gly Ile Ser Asp Val Val

290

295

300

Lys Val Thr Leu Thr His Glu Glu Glu Ala Cys Leu Lys Lys Ser Ala

305

310

315

320

Asp Thr Leu Trp Gly Ile Gln Lys Glu Leu Gln Phe

325

330

<210> 2

<211> 971

<212> DNA

<213> *Saccharomyces cerevisiae*

<400> 2

aagggtagcc tccccataac ataaactcaa taaaatatat agtcttcaac ttgaaaaagg 60
aacaagctca tgc aaagagg tggtagccgc acgccgaaat gcatgcaagt aacctattca 120
aagtaatatc tcatacatgt ttc atgagggg taacaacatg cgactgggtg agcatatgct 180
ccgctgatgt gatgtgcaag ataaacaagc aagacggaaa ctaactcttt cttcatgtaa 240
taaacacacc ccgcgtttat ttacctatct ttaaacttca acaccttata tcataactaa 300
tatttctga gataagcaca ctgcacccat accttcccta aaagcgtagc ttccagtttt 360
tggtaggttc ggcttccctc ccgattccgc ccgctaaacg catatttttg ttgcctgggtg 420
gcatttgcaa aatgcataac ctatgcattt aaaagattat gtatgctctt ctgacttttc 480
gtgtgatgaa gctcgtggaa aaaatgaata atttatgaat ttgagaacaa ttctgtgttg 540
ttacggattt ttactatgga ataattaatc aattgaggat tttatgcaaa tatcgtttga 600
atatttttc gacctttga gtacttttct tcataattgc ataattgt ccgctgccc 660
ttttctgtt agacgggtgc ttgatctact tgctatcgtt caacaccacc ttattttcta 720
actatttttt ttttagctca ttgaaatcag cttatggtga tggcacattt ttgcataaac 780
ctagctgtcc tcgttgaaca taggaaaaaa aaatatatta acaaggctct ttcactctcc 840
ttgcaatcag atttgggttt gttcccttta tttcatatt tctgtcata ttctttctc 900
aattattatt ttctactcat aaccacacgc aaaataaac agtcaaatca atcaagatc 960
ccccaattct c

971

<210> 3

<211> 999

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Modified DNA

coding lactate dehydrogenase

<400> 3

atggctactt tgaagatca attgattcaa aatttgttga aagaagaaca tgtccacaa 60
aataaaatta ctattgttgg tgttgggtgct gttgggatgg ctgtgctat ttctattttg 120
atgaaagatt tggctgatga agttgctttg gttgatgtta tggaagataa attgaaaggt 180
gaaatgatgg atttgaaca tggttctttg tttttgagaa ctccaaaaat tgtttctggt 240
aaagattata atgttactgc taattctaga ttggttatta ttactgctgg tgctagacaa 300
caagaaggtg aatctagatt gaatttggtt caaagaaatg ttaattttt taaatttatt 360
attccaaata ttgttaata ttctccaaat tgtaaatgt tggttgttc taatccagtt 420
gatatattga ctatgttgc ttggaaaatt tctggttttc caaaaaatag agttattggt 480
tctggttgta atttggattc tgctagattt agatatttga tgggtgaaag attgggtgtt 540
catccattgt ctgtcatgg ttggattttg ggtgaacatg gtgattcttc tgtccagtt 600
tggtctggtg ttaatgttgc tgggttttct ttgaaaaatt tgcattcaga attgggtact 660
gatgctgata aagaacaatg gaaagctgtt cataaacaag ttgttgattc tgctatgaa 720
gttattaaat tgaaaggtta tacttcttgg gctattggtt tgtctgttgc tgattggct 780
gaatctatta tgaaaaattt gagaagagtt catccaattt ctactatgat taaaggtttg 840
tatgggtata aagaagatgt tttttgtct gttccatgta ttttgggtca aaatggtatt 900
tctgatgttg ttaaggttac ttgactcat gaagaagaag cttgtttgaa aaaatctgct 960
gatactttgt ggggtattca aaaagaattg caattttaa

999

<210> 4

<211> 1052

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Modified DNA

coding lactate dehydrogenase

<400> 4

acagaattca caatggctac ttgaaagat caatgattc aaaatttgt gaaagaagaa 60
catgtccac aaaataaaat tactattgt ggtgttggtg ctgttggtat ggcttggct 120
attctattt tgatgaaaga ttggctgat gaagttgctt tggttgatgt tatggaagat 180
aaattgaaag gtgaaatgat ggatttgcaa catggttctt tgttttgag aactccaaa 240
attgtttctg gtaaagatta taatgttact gctaattcta gattggttat tattactgct 300
ggtgctagac aacaagaagg tgaatctaga ttgaatttg ttcaagaaa tgtaaatatt 360
ttaaattta ttattccaaa tattgttaa tattctcaa attgtaaat gtgtgttgt 420
tctaaccag ttgatattt gacttatgtt gcttgaaaaa ttctggtt tccaaaaaa 480
agagttaatt gtctggttg taatttggat tctgctagat ttagatatt gatgggtgaa 540
agattgggtg ttcatccatt gtctgtcat ggttgattt tgggtgaaca tgggtattct 600
tctgtccag ttggtctgg tgtaattgt gctggtgtt cttgaaaaa ttgcatcca 660
gaattggga ctgatctga taaagaacaa tggaaagctg ttcataaaca agttgtgat 720
tctgctatg aagtattaa attgaaagg tatactctt gggctattgg ttgtctgt 780
gctgatttg ctgaatctat tatgaaaaa ttgagaagag ttcatcaat ttctactatg 840
attaaagggt tgtatggtat taaagaagat gttttttgt ctgttccatg tatttgggt 900
caaatggta ttctgatgt tgtaaaagt acttgactc atgaagaaga agctgtttg 960
aaaaaatctg ctgatactt gtgggggtatt caaaaagaat tgcaattta ataactcgag 1020
cttggttgaa caggttgcca aggcctaagt ga

1052

<210> 5

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 5

acagaattca caatggctac ttgaaagat caattgattc aaaattgtt gaaagaagaa 60

catgttcac aaaataaaat tactattgtt ggtgttggtg

100

<210> 6

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 6

acagaattca caatggctac

20

<210> 7

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 7

atgataacaa ccacaaccac gacaaccata ccgaacacga taaagataaa actactttct 60
aaaccgacta cttcaacgaa accaactaca ataccttcta

100

<210> 8

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 8

atgataacaa ccacaaccac

20

<210> 9

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 9

tggttgatgt tatggaagat aaattgaaag gtgaaatgat ggatttgcaa catggttctt 60
tgtttttgag aactccaaaa attgtttctg gtaaagatta

100

<210> 10

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 10

tggttgatgt tatggaagat

20

<210> 11

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 11

taacaaagac catttctaatt attacaatga cgattaagat ctaaccaata ataatgacga 60
ccacgatctg ttgttcttc acttagatct aacttaaacc

100

<210> 12

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 12

taacaaagac catttcta a

21

<210> 13

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 13

tgaatctaga ttgaattgg ttcaaagaaa tgtaatat tttaaattta ttattccaaa 60
tattgttaaa tattctcaa attgtaaatt gttggtgtt

100

<210> 14

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 14

tgaatctaga ttgaattgg t

21

<210> 15

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 15

taacatttaa caaccaacaa agattagtc aactataaaa ctgaatacaa cgaacctttt 60
aaagacacaaa aggtttttta tctcaataac caagaccaac

100

<210> 16

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 16

taacatttaa caaccaacaa a

21

<210> 17

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 17

agagttattg gttctggttg taattggat tctgctagat ttagatattt gatgggtgaa 60

agattgggtg ttcattcatt gtctgtcat gttggattt

100

<210> 18

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 18

agagttattg gttctggttg t

21

<210> 19

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 19

cagaacagta ccaacctaaa acccactgt accactaaga agacaaggtc aaaccagacc 60
acaattacaa cgaccacaaa gaaactttt aaacgtaggt 100

<210> 20

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 20

cagaacagta ccaacctaaa a 21

<210> 21

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 21

ctttgaaaaa ttgcatcca gaattgggta ctgatgctga taaagaacaa tggaaagctg 60
ttcataaaca agttgtgat tctgcttatg aagttattaa 100

<210> 22

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 22

ctttgaaaaa ttgcatcca g 21

<210> 23

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 23

agacgaatac ttcaataatt taactttcca atatgaagaa cccgataacc aaacagacaa 60

cgactaaacc gacttagata atacttttta aactcttctc

100

<210> 24

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 24

agacgaatac ttcaataatt t

21

<210> 25

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 25

tatgaaaaat ttgagaagag ttcacccaat ttctactatg attaaagggtt tgtatggat 60

taaagaagat gtttttttgt ctgtccatg tattttgggt

100

<210> 26

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 26

atgaaaaatt tgagaagagt

20

<210> 27

<211> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 27

gacaaggtac ataaaaccca gttttaccat aaagactaca acaatttcaa tgaaactgag 60

tactttctct tgaacaaac ttttttagac gactatgaaa

100

<210> 28

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 28

gacaaggtag ataaaaccca g

21

<210> 29

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 29

aaaaaatctg ctgatacttt gtggggatt caaaaagaat tgcaatttta ataactcgag 60

<210> 30

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 30

aaaaaatctg ctgatacttt g

21

<210> 31

<211> 52

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 31

acgttaaaat tattgagctc gaaccaactt gtgcaacggt tccgaattca ct 52

<210> 32

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 32

acgttaaaat tattgagctc g 21

<210> 33

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 33

atatatggat ccgcgtttat ttacctatct c

31

<210> 34

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 34

atatatgaat tctttgattg attgactgt g

31

<210> 35

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 35

atatatctcg aggccagcta acttcttggt cgac

34

<210> 36

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 36

atatatgaat tctttgattg attgactgt g

31

<210> 37

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 37

tggttgatgt tatggaagat

20

<210> 38

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 38

gacaaggtac ataaaacca g

21

<210> 39

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 39

gtaataaaca caccgccg

19

[Brief explanation of drawings]

[Figure 1]

Figure 1 is a diagram illustrating the Codon Usage method used for *Saccaromyces cerevisiae* and its usage frequency.

[Figure 2]

Figure 2 is a diagram illustrating the homologue analysis results of the base sequence in a bovine-derived LDH and the base sequence in a modified LDH. The top row indicates the base sequence in the bovine-derived LDH while the bottom row indicates the modified base sequence, in which the bases that are different from the base sequence in the bovine-derived LDH are indicated by symbols (A, T, C or G).

[Figure 3]

Figure 3 is a diagram illustrating the primer structure for synthesizing a long-chain DNA by means of PCR used in Embodiment 1 and the synthesis steps (Steps 1 through 4).

[Figure 4]

Figure 4 is a diagram illustrating the plasmid map of the constructed vector pBTrp-PDC1-LDHKCB.

[Figure 5]

Figure 5 is a diagram illustrating a part of the process of constructing the vector shown in Figure 5.

[Figure 6]

Figure 6 is a diagram illustrating another part of the pBTrp-PDC1-LDH construction process.

[Figure 7]

Figure 7 is a diagram illustrating another part of the pBTrp-PDC1-LDH construction process.

[Figure 8]

Figure 8 is a diagram illustrating another part of the process (the last step) of constructing the vector shown in Figure 5.

[Figure 9]

Figure 9 is a diagram illustrating a structural change in the target site of the chromosome DNA in the transformant obtained in one embodiment.

[Figure 10]

Figure 10 is a diagram illustrating the volume percentages of lactic acid and the ethanol produced in one embodiment.

[Figure 11]


Figure 11 is a diagram illustrating the trends in the volume percentages of lactic acid and ethanol produced in one embodiment.

【FIG.1】


	Sec	Tyr	Cys
Phe	UUU 26.0 (146412)	UUA 18.8 (105357)	UGU 8.0 (44797)
	UUC 18.2 (102353)	UAC 14.7 (82477)	UGC 4.7 (26363)
	UUA 26.4 (148212)	UAA 1.0 (5537)	UGA 0.6 (3456)
Leu	UUG 27.1 (152577)	UAG 0.5 (2629)	UGG 10.3 (58084)
	Pro	His	
	CUU 12.2 (68500)	GAU 13.7 (77276)	CGU 6.5 (36513)
	CUC 5.4 (30213)	CAC 7.8 (43867)	CGC 2.6 (14559)
	CUA 13.4 (75415)	CAA 27.5 (154545)	CGA 3.0 (16965)
	CUG 10.4 (58568)	CAG 12.2 (68453)	CGG 1.7 (9606)
		Asn	
Ile	AUU 30.2 (159875)	AAU 36.0 (202449)	AGU 14.2 (79666)
	AUC 17.1 (96127)	AAC 24.9 (140174)	AGC 9.7 (54339)
	AUA 17.8 (100079)	AAA 42.1 (236938)	AGA 21.3 (119693)
Met	AUG 20.9 (117420)	AAG 30.8 (173181)	AGG 9.3 (52060)
	Ala	Asp	Gly
Val	GUU 22.0 (123771)	GAU 21.1 (118604)	Arg 23.9 (134536)
	GUC 11.6 (65195)	GCC 12.6 (70752)	GCG 9.7 (54619)
	GUA 11.8 (66110)	GCA 16.2 (91026)	GGA 10.9 (61498)
	GUG 10.7 (60001)	GCG 6.1 (34530)	GGG 6.0 (33624)
			Glu

[illegible]

【FIG.3】




[FIG.4]




Plasmid name: pBTrp-PDC1-LDHKCB

Plasmid size: 7.13 kb


[FIG.5]




【FIG.6】




[FIG. 7]




【FIG.8】




【FIG.9】



【FIG.10】



【FIG.11】



[Name of Document] Abstract

[Abstract]

[Problems to be Solved]

The present invention provides a transformant into which has been incorporated DNA for coding a foreign protein having lactate dehydrogenase activity and provided with pyruvic acid substrate affinity that equals or exceeds the pyruvic acid substrate affinity of the pyruvate decarboxylase inherent in the host organism. Said transformant can stably mass-produce lactic acid inside a host organism having the pyruvate decarboxylase gene.

[Selected Figure] NONE